

**Drug Facts**

Active ingredient Purpose

Benzocaine 20%.....Oral pain reliever

Uses temporary relief of pain due to minor irritation or injury of the mouth and gums or temporary relief of pain associated with canker sores, dentures or orthodontic appliances

**Warnings**

Do not use

- for more than 7 days unless directed by a dentist or physician
- if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other 'caine' anesthetics

**When using this product**

- do not exceed recommended dosage
- avoid contact with eyes
- localized allergic reactions may occur after prolonged or repeated use

Stop use and ask a doctor if

- sore mouth symptoms do not improve in 7 days
- swelling, rash or fever develops
- irritation, pain, or redness persists or worsens

Keep out of reach of children. If more than used for pain relief is accidentally swallowed, get medical help or contact a Poison Control Center immediately.

**Directions**

Adults and children 2 years and over:

- apply to the affected area with a cotton swab
- use up to 4 times daily or as directed by a dentist or physician
- children under 12 years of age should be supervised in the use of the product
- children under 2 years of age : consult a dentist or doctor

Other information Store at 20 -25°C (68 -77°F)

Inactive ingredients flavor, polyethylene glycol, sodium saccharin

Questions or comments? call toll free 9am to 5pm EST at 1-800-872-8305