

DRUG FACTS

ACTIVE INGREDIENT:

Benzocaine 20%

PURPOSE:

Oral Anesthetic intended for adults and children 2 years of age and older.

USE:

ProJel-20® is a 20% benzocaine topical anesthetic gel packaged in 60 gram jars for use in the operator. The gel is indicated for temporary relief of pain associated with minor irritation or injury in the mouth and gums or of pain due to canker sores, denture sores or orthodontic appliances.

ProJel-20® is available in four flavors: Cherry, Pina Colada, Bubblegum and Mint.

WARNINGS

Methemoglobinemia warning:

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy alert: Do not use on patients with history of allergy due to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

DO NOT USE

- For more than 7 days unless directed by a dentist or physician
- If you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other 'caine' anesthetics
- **For teething**
- **In children under 2 years of age**

WHEN USING THIS PRODUCT

- Do not exceed recommended dosage
- Patient should not ingest or swallow
- Avoid contact with eyes
- Localized allergic reactions may occur after prolonged or repeated use
- Discontinue use if irritation/inflammation or fever and infection develops

STOP USE AND ASK A DOCTOR IF

- Sore mouth symptoms do not improve in 7 days
- Swelling, rash or fever develops
- Irritation, pain, or redness persists or worsens.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

If more than used for pain relief is accidentally swallowed, get medical help or contact a Poison Control Center immediately.

DIRECTIONS:

Apply only amount needed to the oral mucosa to prevent or relieve pain.

Children under 2 years of age: do not use.

OTHER INFORMATION

For professional use only in the practice of dentistry.

Store at 20°C - 25°C (68°F - 77°F). Do not permit to freeze.

HOW SUPPLIED

ProJel-20®, Benzocaine 20% is available in 60 gram jars (NDC 0362-0124-05).

INACTIVE INGREDIENTS:

Flavor, polyethylene glycol, sodium saccharin.

Manufactured by:

SEPTODONT, Inc.

416 S. Taylor Ave

Louisville, CO 80027

800-872-8305 | www.septodontusa.com

Rev. 10/2018

